



CEP57 gene

centrosomal protein 57

Normal Function

The *CEP57* gene provides instructions for making a protein whose function is not completely understood. Within cells, the *CEP57* protein is located in structures called centrosomes. Centrosomes have a role in cell division and the assembly of microtubules. Microtubules are fibers that help cells maintain their shape, assist in the process of cell division, and are essential for the movement (transport) of materials within cells. *CEP57* seems especially important for the organization and stability of specialized microtubules called spindle microtubules, which are important for cell division. Before cells divide, they copy all of their chromosomes. Spindle microtubules, which are produced by centrosomes, attach to the duplicated chromosomes and pull one copy of each to opposite ends of the cell so that each new cell contains one complete set of chromosomes.

The *CEP57* protein is also involved in the transport of certain molecules along microtubules, particularly a protein called fibroblast growth factor 2 (FGF2). FGF2 is an important signaling molecule that helps regulate growth and development of cells and tissues, and its transport inside the cell is important for relaying signals that instruct the cell how to function.

Health Conditions Related to Genetic Changes

Mosaic variegated aneuploidy syndrome


At least three *CEP57* gene mutations have been found to cause mosaic variegated aneuploidy (MVA) syndrome type 2. This condition is characterized by cells with abnormal numbers of chromosomes, a situation known as aneuploidy. Affected individuals grow slowly before and after birth and may have mild intellectual disability and other health problems.

CEP57 gene mutations involved in MVA syndrome type 2 likely reduce the amount of functional *CEP57* protein in cells. It is unclear how these changes lead to the features of the condition. Researchers speculate that impairment of the protein's role in stabilizing spindle microtubules may prevent the normal separation of chromosomes during cell division, leading to aneuploidy. Although they are unsure how *CEP57* gene mutations lead to the other features of MVA syndrome type 2, some suggest that the shortage of functional *CEP57* protein prevents proper transport of FGF2, which may impair cell signaling and lead to problems with growth and development.

Chromosomal Location

Cytogenetic Location: 11q21, which is the long (q) arm of chromosome 11 at position 21

Molecular Location: base pairs 95,790,461 to 95,832,693 on chromosome 11 (Homo sapiens Updated Annotation Release 109.20200522, GRCh38.p13) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- centrosomal protein 57kDa
- centrosomal protein of 57 kDa isoform a
- centrosomal protein of 57 kDa isoform b
- centrosomal protein of 57 kDa isoform c
- FGF2-interacting protein
- KIAA0092
- MVA2
- PIG8
- proliferation-inducing protein 8
- testis-specific protein 57
- translokin
- TSP57

Additional Information & Resources

Educational Resources

- Molecular Biology of the Cell (fourth edition, 2002): Cytoskeletal Machines Perform Both Mitosis and Cytokinesis
https://www.ncbi.nlm.nih.gov/books/NBK26931/#_A3337_
- Molecular Biology of the Cell (fourth edition, 2002): Mitosis
<https://www.ncbi.nlm.nih.gov/books/NBK26934/>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28%28CEP57%5BTIAB%5D%29+OR+%28centrosomal+protein+57%5BTIAB%5D%29%29+OR+%28%28FGF2-interacting+protein%5BTIAB%5D%29+OR+%28KIAA0092%5BTIAB%5D%29+OR+%28MVA2%5BTIAB%5D%29+OR+%28PIG8%5BTIAB%5D%29+OR+%28TSP57%5BTIAB%5D%29+OR+%28centrosomal+protein+57kDa%5BTIAB%5D%29+OR+%28centrosomal+protein+of+57+kDa+isoform+a%5BTIAB%5D%29+OR+%28centrosomal+protein+of+57+kDa+isoform+b%5BTIAB%5D%29+OR+%28centrosomal+protein+of+57+kDa+isoform+c%5BTIAB%5D%29+OR+%28proliferation-inducing+protein+8%5BTIAB%5D%29+OR+%28testis-specific+protein+57%5BTIAB%5D%29+OR+%28translokin%5BTIAB%5D%29%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

Catalog of Genes and Diseases from OMIM

- CENTROSOMAL PROTEIN, 57-KD
<http://omim.org/entry/607951>

Research Resources

- Atlas of Genetics and Cytogenetics in Oncology and Haematology
<http://atlasgeneticsoncology.org/Genes/CEP57ID43008ch11q21.html>
- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=CEP57%5Bgene%5D>
- HGNC Gene Symbol Report
https://www.genenames.org/data/gene-symbol-report/#!/hgnc_id/HGNC:30794
- Monarch Initiative
<https://monarchinitiative.org/gene/NCBIGene:9702>
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/9702>
- UniProt
<https://www.uniprot.org/uniprot/Q86XR8>

Sources for This Summary

- OMIM: CENTROSOMAL PROTEIN, 57-KD
<http://omim.org/entry/607951>
- Meunier S, Navarro MG, Bossard C, Laurell H, Touriol C, Lacazette E, Prats H. Pivotal role of translokin/CEP57 in the unconventional secretion versus nuclear translocation of FGF2. *Traffic*. 2009 Dec;10(12):1765-72. doi: 10.1111/j.1600-0854.2009.00985.x. Epub 2009 Sep 14.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/19804566>

- Pinson L, Mannini L, Willems M, Cucco F, Sirvent N, Frebourg T, Quarantotti V, Collet C, Schneider A, Sarda P, Geneviève D, Puechberty J, Lefort G, Musio A. CEP57 mutation in a girl with mosaic variegated aneuploidy syndrome. *Am J Med Genet A*. 2014 Jan;164A(1):177-81. doi: 10.1002/ajmg.a.36166. Epub 2013 Nov 20.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/24259107>
- Snape K, Hanks S, Ruark E, Barros-Núñez P, Elliott A, Murray A, Lane AH, Shannon N, Callier P, Chitayat D, Clayton-Smith J, Fitzpatrick DR, Gisselsson D, Jacquemont S, Asakura-Hay K, Micale MA, Tolmie J, Turnpenny PD, Wright M, Douglas J, Rahman N. Mutations in CEP57 cause mosaic variegated aneuploidy syndrome. *Nat Genet*. 2011 Jun;43(6):527-9. doi: 10.1038/ng.822. Epub 2011 May 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/21552266>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3508359/>
- Wu Q, He R, Zhou H, Yu AC, Zhang B, Teng J, Chen J. Cep57, a NEDD1-binding pericentriolar material component, is essential for spindle pole integrity. *Cell Res*. 2012 Sep;22(9):1390-401. doi: 10.1038/cr.2012.61. Epub 2012 Apr 17.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/22508265>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC3434346/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/CEP57>

Reviewed: July 2017
Published: June 23, 2020

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services